

UCLA

UCLA Previously Published Works

Title

Host cell invasion and virulence mediated by Candida albicans Ssa1.

Permalink

<https://escholarship.org/uc/item/54x417fk>

Journal

PLoS pathogens, 6(11)

ISSN

1553-7366

Authors

Sun, Jianing N
Solis, Norma V
Phan, Quynh T
et al.

Publication Date

2010-11-01

DOI

10.1371/journal.ppat.1001181

Peer reviewed

Host Cell Invasion and Virulence Mediated by *Candida albicans* Ssa1

Jianing N. Sun¹, Norma V. Solis², Quynh T. Phan², Jashanjot S. Bajwa¹, Helena Kashleva³, Angela Thompson³, Yaoping Liu², Anna Dongari-Bagtzoglou³, Mira Edgerton^{1*}, Scott G. Filler^{2,4*}

1 Department of Oral Biology, School of Dental Medicine, State University of New York at Buffalo, Buffalo, New York, United States of America, **2** Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, California, United States of America, **3** Division of Periodontology, School of Dental Medicine, University of Connecticut, Farmington, Connecticut, United States of America, **4** The David Geffen School of Medicine at UCLA, Los Angeles, California, United States of America

Abstract

Candida albicans Ssa1 and Ssa2 are members of the HSP70 family of heat shock proteins that are expressed on the cell surface and function as receptors for antimicrobial peptides such as histatins. We investigated the role of Ssa1 and Ssa2 in mediating pathogenic host cell interactions and virulence. A *C. albicans* *ssa1Δ/Δ* mutant had attenuated virulence in murine models of disseminated and oropharyngeal candidiasis, whereas an *ssa2Δ/Δ* mutant did not. *In vitro* studies revealed that the *ssa1Δ/Δ* mutant caused markedly less damage to endothelial cells and oral epithelial cell lines. Also, the *ssa1Δ/Δ* mutant had defective binding to endothelial cell N-cadherin and epithelial cell E-cadherin, receptors that mediate host cell endocytosis of *C. albicans*. As a result, this mutant had impaired capacity to induce its own endocytosis by endothelial cells and oral epithelial cells. Latex beads coated with recombinant Ssa1 were avidly endocytosed by both endothelial cells and oral epithelial cells, demonstrating that Ssa1 is sufficient to induce host cell endocytosis. These results indicate that Ssa1 is a novel invasin that binds to host cell cadherins, induces host cell endocytosis, and is critical for *C. albicans* to cause maximal damage to host cells and induce disseminated and oropharyngeal disease.

Citation: Sun JN, Solis NV, Phan QT, Bajwa JS, Kashleva H, et al. (2010) Host Cell Invasion and Virulence Mediated by *Candida albicans* Ssa1. PLoS Pathog 6(11): e1001181. doi:10.1371/journal.ppat.1001181

Editor: Stuart M. Levitz, University of Massachusetts Medical School, United States of America

Received: March 10, 2010; **Accepted:** October 6, 2010; **Published:** November 11, 2010

Copyright: © 2010 Sun et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported in part by grants R01DE010641, R01AI054928, R01DE017088, R01DE13986, and M01RR00425 from the National Institutes of Health, USA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: SGF holds equity in NovaDigm Therapeutics.

* E-mail: edgerto@buffalo.edu (ME); sfiller@ucla.edu (SGF)

Introduction

The fungus, *Candida albicans* is a significant human pathogen. In hospitalized patients, this organism disseminates hematogenously and infects virtually all organs. Even with currently available therapy, bloodstream infections with *C. albicans* are associated with a 37% mortality [1]. *C. albicans* is also part of the normal oral flora and it usually grows as a harmless commensal. However, when local or systemic host defense mechanisms are impaired, this organism can proliferate and cause debilitating oropharyngeal candidiasis.

To persist within the human host and cause disease, *C. albicans* must be able to adhere to and invade host cells or tissues while resisting the stress caused by host-derived reactive oxygen intermediates and antimicrobial peptides [2–5]. In other organisms, heat shock proteins play an important role in each of these activities. For example, some heat shock proteins are expressed on the cell surface of microorganisms, where they function as adhesins [6–9]. Also, in some bacteria and parasites, members of the Hsp70 and Hsp100 family of heat shock proteins are required for resistance to host-induced stress [10–12].

Ssa1 and Ssa2 are the only two members of the Hsp70 family in *C. albicans*, and both proteins are expressed on the cell surface of yeast and hyphae [13,14]. Previously, we found that histatin 5, one of the main antimicrobial proteins found in saliva, binds with high affinity to Ssa2 and with lower affinity to Ssa1. After histatin 5 is bound to Ssa proteins, it is transported into the cytoplasm, where it

kills the fungal cell [15,16]. *C. albicans* Ssa1 and Ssa2 are also required for maximal fungicidal activity of human β defensins 2 and 3 [17].

As reported here, we investigated the roles of Ssa1 and Ssa2 in *C. albicans* virulence in murine models of hematogenously disseminated and oropharyngeal candidiasis. We found that Ssa1, but not Ssa2 is essential for normal virulence in both models. Through *in vitro* experiments, we discovered that surface-expressed Ssa1 likely contributes to virulence by acting as an invasin and directly mediating *C. albicans* invasion of both endothelial and oral epithelial cells *in vitro*.

Results

Ssa1 is required for maximal virulence in a mouse model of hematogenously disseminated candidiasis

To evaluate the contribution of Ssa1 and Ssa2 to *C. albicans* pathogenicity, the virulence of *ssa1Δ/Δ* and *ssa2Δ/Δ* null mutants was tested in a murine model of disseminated candidiasis. Over the 21 day observation period, none of the mice infected intravenously with the *ssa1Δ/Δ* mutant died (Figure 1A). In contrast, the median survival of mice infected with the wild-type and *ssa1Δ/Δ::SSA1Δ/Δ* complemented strain was 7 to 8 days. Ssa2 did not appear to influence virulence in this model because the survival of mice infected with the *ssa2Δ/Δ* null mutant was similar to that of mice infected with the wild-type strain ($p = 0.13$).

Author Summary

The fungus *Candida albicans* can proliferate in the mouth, causing oropharyngeal candidiasis. In other patients, it can enter the bloodstream and spread throughout the body, resulting in hematogenously disseminated candidiasis. Fungal invasion of host cells is a key feature of both types of infection. One mechanism by which *C. albicans* invades both the epithelial cell lining of the oropharynx and the endothelial cell lining of the blood vessels is by inducing its own uptake. This uptake is induced in part by the binding of the *C. albicans* invasin Als3 to host cell proteins, which include N- and E-cadherin. Here we show that *C. albicans* Ssa1, a member of the 70 kDa heat shock protein family, is expressed on the surface of *C. albicans* where it functions as an invasin. The key role of Ssa1 in host cell invasion is illustrated by the reduced capacity of an *ssa1Δ/Δ* null mutant to induce its own uptake by epithelial and endothelial cells in vitro, and by the significantly attenuated virulence of this mutant in mouse models of oropharyngeal candidiasis and disseminated candidiasis. Thus, Ssa1 is the second identified invasin of *C. albicans*.

Next, we compared the organ fungal burden at various time points in mice infected with the different strains. After 8 h of infection, the kidney and brain fungal burdens of mice inoculated with the *ssa1Δ/Δ* mutant were significantly lower than mice infected with either the wild-type or *ssa1Δ/Δ::SSA1Δ/Δ* complemented strains (Figures 1B–D). After 1 and 2 days of infection, the kidney and brain fungal burdens of mice challenged with the *ssa1Δ/Δ* mutant either declined or were stable. In contrast, the fungal burden of these organs steadily increased in mice infected with either the wild-type or *ssa1Δ/Δ::SSA1Δ/Δ* complemented strains. The fungal burdens of the livers of mice infected with the *ssa1Δ/Δ* mutant were also significantly lower than those of the control mice after 1 and 2 days of infection. These results indicate that *SSA1* is required for normal levels of *C. albicans* infection in the kidney and brain as early as 8 h after inoculation. *SSA1* also appears to be necessary for the organism to persist in the tissues at later time points.

To investigate why mice infected with the *ssa1Δ/Δ* mutant had no mortality over the course of the infection, we determined their organ fungal burden after 4, 7 and 14 days post-inoculation. We were not able to analyze the organ fungal burden of mice infected with either the wild-type strain or the *ssa1Δ/Δ::SSA1Δ/Δ* complemented strain at these later time points because a significant percentage of these mice had died. We found that mice infected with the *ssa1Δ/Δ* mutant progressively cleared the infection so that their organs became sterile by 14 days (Figure 1B–D and data not shown). This clearance provides an explanation for the prolonged survival of mice infected with the *ssa1Δ/Δ* mutant.

The low fungal burden in the kidneys of mice infected with the *ssa1Δ/Δ* mutant was verified by histopathology. After 1 day of infection, the kidneys of these mice contained very small foci of organisms. These foci usually consisted of a single fungal element surrounded by a few inflammatory cells (Figure 1E). Interestingly, many of the *ssa1Δ/Δ* cells were located in the glomeruli. The kidneys from mice infected with either the wild-type or *ssa1Δ/Δ::SSA1Δ/Δ* complemented strain contained the expected micro-abscesses, which contained numerous neutrophils surrounding multiple *C. albicans* filaments. It was not possible to determine whether the filaments of the various *C. albicans* strains were true hyphae or pseudohyphae in these histopathologic specimens. However, the filaments of the *ssa1Δ/Δ* mutant appeared to be

similar in length to those of the wild-type and *ssa1Δ/Δ::SSA1Δ/Δ* complemented strain. Also, we have previously found that the *ssa1Δ/Δ* mutant forms hyphae similar to the wild-type strain in the presence of serum in vitro [16]. Therefore, the attenuated virulence of the *ssa1Δ/Δ* mutant was unlikely due to a defect in filamentation.

During the construction of the *ssa1Δ/Δ* mutant, *URA3* was used as the selectable marker. The chromosomal locus at which *URA3* is integrated can sometimes influence the virulence of *C. albicans* mutants in the mouse model of disseminated candidiasis [18–20]. In the *ssa1Δ/Δ* mutant, *URA3* was integrated at the *SSA2* locus [16]. However, in the *ssa1Δ/Δ::SSA1* complemented strain, *URA3* was integrated at the *RPS10* locus, which is known to result in normal activity of the *URA3* gene product, orotidine 5'-monophosphate decarboxylase [21]. To verify that the observed attenuated virulence of the *ssa1Δ/Δ* mutant was due to the absence of *SSA1* and not the result of the chromosomal locus of *URA3*, we tested the virulence of a second *ssa1Δ/Δ* strain in which *URA3* was integrated at the *RPS10* locus. As expected, all mice infected intravenously with this mutant survived (Supplemental Figure S1A), thus confirming that *SSA1* is required for the normal virulence of *C. albicans*.

The *ssa1Δ/Δ* mutant had attenuated virulence in the mouse model of oropharyngeal candidiasis

Next, we investigated the contributions of *SSA1* and *SSA2* to virulence in a mouse model of oropharyngeal candidiasis. There was a trend towards reduced oral fungal burden in mice infected with the *ssa1Δ/Δ* mutant after 1 day of infection, but this difference was only significant when compared to the *ssa1Δ/Δ* complemented strain ($p=0.04$), but not the wild-type strain ($p=0.13$) (Figure 2A). However, after 2 and 5 days of infection, mice infected with the *ssa1Δ/Δ* mutant had markedly lower oral fungal burdens compared to mice infected with either the wild-type or *ssa1Δ/Δ::SSA1* complemented strain. In contrast, the oral fungal burden of mice infected with the *ssa2Δ/Δ* mutant was similar to that of mice infected with the wild-type strain ($p=0.57$) (Figure 2B). Therefore, Ssa1 is required for maximal virulence during oropharyngeal candidiasis, but Ssa2 is dispensable for virulence during this infection.

The results of histopathologic examination of the tongues of the mice infected with the various strains verified the reduced virulence of the *ssa1Δ/Δ* mutant. The tongues of mice infected with the *ssa1Δ/Δ* mutant had relatively small lesions that contained fewer fungal cells and neutrophils than the lesions of mice infected with either the wild-type or the *ssa1Δ/Δ::SSA1* complemented strain (Figure 2C). Importantly, the filaments of the *ssa1Δ/Δ* mutant were similar in length to those of the wild-type strain. Collectively, these results indicate that Ssa1 is necessary for maximal *C. albicans* virulence during both hematogenously disseminated and oropharyngeal candidiasis.

Ssa1 is required for *C. albicans* to cause maximal damage to both endothelial and oral epithelial cells in vitro

During hematogenously disseminated candidiasis, blood-borne *C. albicans* cells must adhere to and penetrate the endothelial cell lining of the blood vessels to invade the deep tissues [3]. *C. albicans* also adheres to and invades oral epithelial cells during oropharyngeal candidiasis [22–24]. After the organism invades endothelial or oral epithelial cells *in vitro*, it damages these cells. Moreover, *C. albicans* mutants with impaired capacity to cause host cell damage *in vitro* frequently have attenuated virulence in mice [25–27]. Therefore, we used a ^{51}Cr release assay to determine the capacity of the *ssa1Δ/Δ* mutant to damage monolayers of human




Figure 1. Ssa1 is required for normal virulence during hematogenously disseminated candidiasis. Mice were inoculated via the tail vein with 5×10^5 yeast-phase cells of the indicated strains of *C. albicans*. (A) Survival over 21 days ($n = 10$ mice per strain of *C. albicans*). $*P < 0.0001$ compared to mice infected with either the wild-type or *ssa1Δ/Δ::SSA1* complemented strain. (B–D) Fungal burden of the kidneys (B), brains (C), and livers (D) of mice infected with the wild-type, *ssa1Δ/Δ* and *ssa1Δ/Δ::SSA1* complemented strains at the indicated time points. For each organ, the lower bound of the y-axis scale indicates the limits of detection. Results from the 1 day time point are the median \pm interquartile range of two experiments with a total of 12 mice per strain of *C. albicans*. Results from the other time points are the median \pm interquartile range of one experiment with 7 mice per strain $*P < 0.02$ compared to mice infected with either the wild-type or *ssa1Δ/Δ::SSA1* complemented strain. (E) Periodic acid Schiff stained thin sections of kidneys after 1 day of infection with the indicated strains of *C. albicans*. Arrows indicate *C. albicans* filaments in the tissues.

doi:10.1371/journal.ppat.1001181.g001

umbilical vein endothelial cells and the FaDu oral epithelial cell line. In this assay, the host cells are incubated with ^{51}Cr , which is taken up by the cells and binds to cytoplasmic proteins. When the host cells are damaged, there is loss of membrane integrity and the labeled proteins leak out of the cells into the medium. The amount of ^{51}Cr that is released is proportional to the extent of cellular damage [26–31]. We found that the *ssa1Δ/Δ* mutant caused 50% less damage to endothelial cells and 89% less damage to epithelial cells compared to the wild-type strain (Figure 3). Complementing the *ssa1Δ/Δ* mutant with a wild-type copy of *SSA1* restored its capacity to damage these cells. As predicted by the virulence data, the *ssa2Δ/Δ* mutant caused the same extent of damage to the endothelial and epithelial cells as did the wild-type strain. All

strains of *C. albicans* formed true hyphae that were of similar length on both endothelial and epithelial cells (data not shown). Therefore, the host cell damage defects of the *ssa1Δ/Δ* mutant were not due to impaired hyphal formation [27,32,33]. These results indicate that Ssa1 is necessary for *C. albicans* to cause maximal damage to endothelial and oral epithelial cells *in vitro*. This defective capacity to damage host cells likely contributed to the attenuated virulence of the *ssa1Δ/Δ* mutant.

Although endothelial cells grow as a monolayer *in vivo*, epithelial cells grow in multiple, stratified layers in the oropharynx. Therefore, we tested the interactions of the *ssa1Δ/Δ* and *ssa2Δ/Δ* mutants with oral epithelial cells in a three-dimensional culture model, which more closely mimics the normal human oral mucosa and




Figure 2. Attenuated virulence of the *ssa1Δ/Δ* mutant during oropharyngeal candidiasis. Mice were immunosuppressed with cortisone acetate and then orally inoculated with yeast-phase cells of the indicated strains of *C. albicans*. After 1, 2, and 5 days of infection, the mice were sacrificed and the tongues and attached tissues were excised. (A) Oral fungal burden of mice infected with the wild-type, *ssa1Δ/Δ*, and *ssa1Δ/Δ::SSA1* complemented strains for the indicated times. Results from the 1 and 2 day time points are the median \pm the interquartile range of one experiment with 7 mice per strain of *C. albicans*. Results from the 5 day time point are the median \pm the interquartile range of two experiments with a total of 12 mice per strain. * $P \leq 0.01$ compared to both the wild-type and *ssa1Δ/Δ::SSA1* strains. (B) Oral fungal burden of mice infected with the wild-type, *ssa2Δ/Δ*, and *ssa2Δ/Δ::SSA2* complemented strain for 5 days. Results are the median \pm the interquartile range of one experiment with 7 mice per strain of *C. albicans*. (C) Periodic acid-Schiff stained thin sections of the tongues of mice infected with the indicated strains for 5 days. Arrows indicate the *C. albicans* filaments.
doi:10.1371/journal.ppat.1001181.g002

submucosa [34]. Consistent with the intraoral animal findings, the *ssa1Δ/Δ* mutant grew only in small foci and caused little visible epithelial cell damage, even after 2 days of infection (Figure 4A). There was only slight cellular edema in the epithelium, which otherwise maintained its barrier function and prevented the organisms from crossing the basal layer of cells into the collagen gel. In contrast, the *ssa1Δ/Δ::SSA1* complemented strain formed a thick biofilm on the epithelial cells, which resulted in degradation of most of the epithelial layers and subsequent submucosal invasion. As predicted by our previous results, the *ssa2Δ/Δ* strain induced extensive epithelial destruction and invaded into the submucosal compartment similarly to the *ssa1Δ/Δ::SSA1* complemented strain.

To quantify the extent of epithelial cell damage caused by the various strains of *C. albicans* in the 3-dimensional model, we measured the leakage of lactate dehydrogenase (LDH) into the medium. The *ssa1Δ/Δ* mutant induced less LDH release than did the *ssa1Δ/Δ::SSA1* complemented or *ssa2Δ/Δ* strains (Figure 4B). These results provide further support for the importance of Ssa1 in the capacity of *C. albicans* to damage oral epithelial cells.

Ssa1 governs the capacity of *C. albicans* to bind to cadherins and invade endothelial and oral epithelial cells

To cause maximal damage endothelial or oral epithelial cells, *C. albicans* must adhere to and then invade these host cells [27,30,31]. One mechanism by which *C. albicans* invades endothelial cells and oral epithelial cells is by inducing its own endocytosis [27,31,33,35]. Therefore, we used our standard differential fluorescent assay to determine if the *ssa1Δ/Δ* mutant had a defect in its capacity to adhere to and/or be endocytosed by endothelial and FaDu epithelial cells *in vitro*. Compared to the wild-type strain, the *ssa1Δ/Δ* mutant was endocytosed poorly by both endothelial cells and oral epithelial cells (Figure 5A). The *ssa1Δ/Δ::SSA1* complemented strain and the *ssa2Δ/Δ* mutant interacted with both the endothelial and epithelial cells similarly to the wild-type strain. In addition, endocytosis defects of the *ssa1Δ/Δ* mutant persisted when *URA3* was integrated at the *RPS10* locus (Supplemental Figure S1B). Therefore, Ssa1, but not Ssa2, is required for *C. albicans* to invade endothelial cells and oral epithelial cells *in vitro*.

C. albicans induces its own endocytosis by endothelial cells *in vitro* by binding to N-cadherin on the endothelial cell surface [36]. The corresponding receptor for *C. albicans* on oral epithelial cells is E-cadherin [37]. To determine the mechanism of host cell invasion




Figure 3. Ssa1 is necessary for *C. albicans* to cause maximal damage to endothelial cells and an oral epithelial cell line. Endothelial cells and the FaDu oral epithelial cell line were incubated with the indicated strains of *C. albicans* for 3 h, after which the extent of host cell damage was determined using a ⁵¹Cr release assay. Results are the mean \pm SD of three independent experiments, each performed in triplicate. * $P < 0.001$ compared with the wild-type and *ssa1Δ/Δ::SSA1* complemented strains.
doi:10.1371/journal.ppat.1001181.g003




Figure 4. The *ssa1Δ/Δ* mutant has impaired capacity to damage a three-dimensional organotypic model of the oral mucosa. The three-dimensional model of the oral mucosa was infected by adding 10^5 yeast-phase cells of the indicated *C. albicans* strains to the apical surface. (A) Histopathology after 2 days of infection. Thin sections were stained with periodic acid-Schiff. The results are representative of one of three experiments. (B) Epithelial cell damage by the indicated strains was quantified by the release of lactate dehydrogenase (LDH) into the medium. Results are the mean \pm SD of 2 experiments. doi:10.1371/journal.ppat.1001181.g004

defects of the *ssa1Δ/Δ* mutant, we examined its capacity to bind to N-cadherin and E-cadherin in cell membrane extracts from endothelial cells and epithelial cells, respectively. Hyphae of the *ssa1Δ/Δ* mutant bound poorly to N-cadherin and E-cadherin relative to both the wild-type and *ssa1Δ/Δ::SSA1* complemented strain, (Figure 5B). This reduced binding to cadherins likely contributed to the impaired host cell invasion and subsequent reduced host cell damage of the *ssa1Δ/Δ* mutant.

Ssa1 can directly mediate adherence to and endocytosis by host cells

Ssa1 is expressed either on or near the cell surface of both hyphae and yeast phase *C. albicans* [13]. Therefore, we investigated the possibility that Ssa1 by itself could mediate pathogenic host cell interactions. Latex beads were coated with recombinant Ssa1 (rSsa1), rSsa2, or bovine serum albumin (BSA), after which their endocytosis by host cells was measured. Both endothelial cells and epithelial cells endocytosed significantly more beads coated with either rSsa1 or rSsa2 compared to the control BSA-coated beads (Figure 6). These results suggest that both Ssa1 and Ssa2 can directly induce endocytosis by endothelial and epithelial cells.

To verify that Ssa1 was expressed on the surface of *C. albicans* hyphae that were interacting with host cells, we infected FaDu epithelial cells with a strain of *C. albicans* that expressed an Ssa1-GFP fusion protein. After a 90 min incubation, the cells were fixed, stained with an Alexa 594-conjugated anti-*C. albicans* antibody to label the cell surface, and then imaged by confocal microscopy. We found that Ssa1-GFP fluorescence was strongest at the periphery of the hyphae (Figure 7). Importantly, this fluorescence co-localized with that of the fluorescently-labeled anti-*C. albicans* antibody, indicating that Ssa1 was located in the outermost part of the cell

wall. Interestingly, when a similar experiment was performed using organisms grown as yeast cells in suspension, Ssa1 was located slightly internal to the cell surface (Figure S2). These results suggest that the location of Ssa1 is dependent on the morphology and culture conditions of the organism. Importantly, when *C. albicans* hyphae are in contact with host cells, Ssa1 is likely expressed on the cell surface where it can interact with host cell receptors.

Ssa1 and Als3 function in the same pathway(s) to induce endocytosis

One *C. albicans* ligand that binds to N-cadherin and E-cadherin is Als3, and *als3Δ/Δ* mutants have significantly reduced adherence to and invasion of both endothelial cells and oral epithelial cells in vitro [37–39]. Because heat shock proteins can potentially influence the expression of other proteins on the surface of *C. albicans*, we investigated the possibility that the host cell interaction defects of the *ssa1Δ/Δ* mutant were due to decreased surface expression of Als3. Flow cytometric analysis of hyphae stained with a polyclonal anti-Als3 antibody demonstrated that the *ssa1Δ/Δ* mutant expressed a similar amount of Als3 on its surface as did the wild-type and *ssa1Δ/Δ::SSA1* complemented strains (Figure 8A). Furthermore, using indirect immunofluorescence with the anti-Als3 antibody, we determined that Als3 was distributed along the entire length of *ssa1Δ/Δ* mutant hyphae, similar to the wild-type and *ssa1Δ/Δ::SSA1* complemented strains (Figure 8B). Thus, the host cell interaction defects of the *ssa1Δ/Δ* mutant are unlikely to be due to abnormal amount or location of Als3 on the cell surface.

To further investigate possible interactions between Ssa1 and Als3, we constructed an *ssa1Δ/Δ als3Δ/Δ* double mutant and compared its host cell interactions with those of *ssa1Δ/Δ* and *als3Δ/Δ* single mutants. When a 90 min incubation period was used, the host cell interaction defects of the both the *ssa1Δ/Δ* and *als3Δ/Δ* mutants were so large that it was not possible to detect any further reduction in endocytosis of the *ssa1Δ/Δ als3Δ/Δ* double mutant (Figure 5 and data not shown). Therefore, we increased the incubation period to 150 min. We found that at this time point, the *als3Δ/Δ* single mutant had a greater defect in inducing endocytosis by endothelial and epithelial cells than did the *ssa1Δ/Δ* single mutant (Figure 8C). However, the endocytosis defect of the *ssa1Δ/Δ als3Δ/Δ* double mutant was similar to that of the *als3Δ/Δ* single mutant. Collectively, these results indicate that Ssa1 and Als3 function in the same pathway(s) to induce endocytosis, perhaps by binding to the same host cell surface proteins and/or forming part of the same multiprotein complex.

SSA1 is expressed at higher levels than SSA2

It seemed paradoxical that Ssa2 was sufficient to induce endocytosis by host cells, yet the *ssa2Δ/Δ* mutant was endocytosed similarly to the wild-type strain. One potential explanation for these results is that SSA1 is expressed at a higher level than SSA2. Previously, we had found that, in yeast-phase *C. albicans*, the SSA1 mRNA transcript levels were significantly greater than those of SSA2, and the fungal cell wall contained 4- to 5- fold more Ssa1 than Ssa2 [16]. To determine if SSA1 was expressed greater than SSA2 in hyphae that were in contact with host cells, we infected FaDu oral epithelial cells with the various *C. albicans* strains and allowed them to germinate. SSA1 and SSA2 mRNA expression in these organisms was measured by real-time PCR. We found that in the wild-type strain, SSA1 was expressed 8-fold higher than SSA2 (Figure 9). As expected, SSA1 mRNA was not detectable in the *ssa1Δ/Δ* mutant. Importantly, SSA2 expression in this mutant was similar to that of the wild-type strain, indicating that SSA2 was not up-regulated in compensation for the absence of SSA1. Therefore, even though Ssa2 is capable of stimulating host cell




Figure 5. Ssa1 is necessary for *C. albicans* to adhere to and be endocytosed by endothelial cells and oral epithelial cells. (A) Adherence and endocytosis assay. The indicated strains of *C. albicans* were incubated with endothelial cells or FaDu oral epithelial cells for 90 min, after which the number of endocytosed and cell-associated (endocytosed plus adherent) organisms was determined by a differential fluorescent assay. Results are the mean \pm SD of three independent experiments, each performed in triplicate. * $P < 0.005$ compared to the wild-type and *ssa1Δ/Δ::SSA1* complemented strains; † $p < 0.03$ compared to the wild-type and *ssa1Δ/Δ::SSA1* complemented strains. (B) The capacity of the indicated strains to bind to N-cadherin in extracts of endothelial cell membrane proteins and E-cadherin in extracts of FaDu oral epithelial cell membrane proteins was determined using a whole-cell affinity purification approach. The immunoblot of endothelial cell proteins eluted from hyphae of the indicated strains was developed with an anti-N-cadherin antibody. The immunoblot of epithelial cell proteins eluted from hyphae of the indicated strains was developed with an anti-E-cadherin antibody. doi:10.1371/journal.ppat.1001181.g005

endocytosis, the effects of *SSA2* deletion are likely masked by the high level expression of *SSA1*.

Ssa1 is not required for secretion of phospholipases and proteases

C. albicans secretes phospholipases and aspartyl proteases, both of which contribute to the virulence of this organism, probably by damaging host cells [35,40–46]. Heat shock proteins can influence protein secretion [47,48]. Therefore, we investigated the possibility

that the attenuated damage of endothelial and epithelial cells caused by the *ssa1Δ/Δ* mutant was due to reduced secretion of phospholipases or proteases. We grew the various strain on either egg yolk agar or BSA agar and measured the size of the zones of precipitation or clearance around the colonies to screen for total extracellular phospholipase and protease activity, respectively [43,49]. The *ssa1Δ/Δ* mutant was similar to the wild-type and *ssa1Δ/Δ::SSA1* complemented strains in these assays (Figure 10). These data suggest that the reduced virulence of the *ssa1Δ/Δ*




Figure 6. Ssa1 is sufficient to induce endocytosis by host cells. Latex beads were coated with bovine serum albumin (BSA), recombinant Ssa1 (rSsa1), or rSsa2. They were incubated for 45 min with endothelial cells and FaDu oral epithelial cells, after which the number of endocytosed and cell-associated beads was determined. Data are expressed as a percentage of the results with beads coated with BSA and are the mean \pm SD of 3 experiments, each performed in triplicate. * $P < 0.01$ compared to beads coated with BSA. doi:10.1371/journal.ppat.1001181.g006




Figure 7. Ssa1 is located on the cell surface of *C. albicans* hyphae during epithelial cell interaction. FaDu oral epithelial cells were infected with *C. albicans* expressing an Ssa1-GFP fusion protein for 90 min, after which the cells were fixed and stained with an Alexa 594-conjugated anti-*C. albicans* polyclonal antibody to label the cell surface. (A) Confocal microscopic images of Ssa1-GFP (1) and fluorescent-labeled anti-*C. albicans* antibody (2). The merged image is shown in (3) and the regions in yellow indicate areas of co-localization between Ssa1-GFP and the fluorescent-labeled anti-*C. albicans* antibody. (B) Graphs of fluorescent intensity at different cross sections of the hypha in panel (3). The green lines indicate the fluorescent intensity of the Ssa1-GFP and the red lines indicate the fluorescent intensity of the fluorescent-labeled anti-*C. albicans* antibody. The letters (a-c) correspond to those in panel (3) and indicate the locations along the hypha at which the fluorescent intensity was measured. doi:10.1371/journal.ppat.1001181.g007

mutant was unlikely the result of diminished secretion of phospholipases or proteases.

C. albicans resistance to stress is independent of Ssa1 and Ssa2

Heat shock proteins are important for some microorganisms to resist the stressful conditions they encounter in the host [10–12]. Therefore, increased susceptibility to host-induced stress could have contributed to the reduced virulence of the *ssa1Δ/Δ* mutant. To evaluate this possibility, we tested the susceptibility of the *ssa1Δ/Δ* and *ssa2Δ/Δ* mutants to various stressors. Both the *ssa1Δ/Δ* and *ssa2Δ/Δ* mutants had wild-type susceptibility to oxidant (menadione and H₂O₂), cell wall (Calcofluor white), osmotic (NaCl), and plasma membrane (SDS) stress (Figure 11A). To verify that deletion of *SSA1* or *SSA2* did not affect stress resistance, we tested these stressors at higher concentrations. The growth of the *ssa1Δ/Δ* and *ssa2Δ/Δ* mutants was inhibited similarly to the wild-type strain under these conditions (data not shown). In addition, all strains had comparable susceptibility to nitric oxide (data not shown). We also tested the susceptibility of the *ssa1Δ/Δ* mutant to damage by the human neutrophil-like HL-60 cell line. This mutant was not more susceptible than the wild-type and *ssa1Δ/Δ::SSA1* complemented strains ($p \geq 0.12$) (Figure 11B). Therefore, the reduced virulence of the *ssa1Δ/Δ* mutant was unlikely to be due to increased susceptibility to host induced stress.

Discussion

Heat shock proteins play diverse roles in the pathogenicity of many microorganisms, including bacteria, protozoa, and fungi [6–12]. However, the contribution of members of the HSP70 family of heat shock proteins to *C. albicans* virulence has not been reported previously. Our results with the *ssa1Δ/Δ* mutant demonstrate that Ssa1 is required for maximal *C. albicans* virulence during both hematogenously disseminated and oropharyngeal candidiasis in mice.

The data from our *in vitro* studies indicated that the *ssa1Δ/Δ* mutant was defective in its capacity to adhere to, invade, and damage both endothelial cells and oral epithelial cells. The invasion defect of this mutant was likely due in part to its impaired capacity to bind to endothelial cell N-cadherin and epithelial cell E-cadherin, receptors that can mediate the endocytosis of *C. albicans* [36,37,39]. Other *C. albicans* mutants that are defective in invading and damaging endothelial cells or oral epithelial cells frequently have attenuated virulence in mouse models of candidiasis [25,26,32,50]. Therefore, it is highly probable that the host cell interaction defects of the *ssa1Δ/Δ* mutant contributed to its attenuated virulence in mice.

The experiments with latex beads coated with Ssa1 or Ssa2 demonstrated that either protein can induce endocytosis by both endothelial cells and oral epithelial cells. However, in intact organisms, Ssa1 is likely more important than Ssa2 for inducing host cell endocytosis because the transcript levels of *SSA1* are much greater. In addition, using GFP-labeled Ssa1, we verified that Ssa1 is expressed on the surface of hyphae that are in contact with epithelial cells. Thus, Ssa1 is located where it can bind to host cell receptors. Collectively, these results indicate that Ssa1 can function as an invasin and induce host cells to endocytose the organism. Members of the Hsp70 family have also been found to mediate the host cell adherence of bacteria such as *Helicobacter pylori*, *Haemophilus influenza*, and *Mycobacterium avium* [6–8]. In addition, Hsp60 is expressed on the surface of *Histoplasma capsulatum* and is the major ligand for CD11b/CD18 on macrophages [9]. Orthologs of Ssa1 and Ssa2 are also present in species of *Candida* other than *C. albicans*. Whether these orthologs also mediate pathogenic host cell interactions remains to be determined.

Ssa1 is functionally similar to the *C. albicans* invasin Als3, which also binds to cadherins on the surface of endothelial cells and epithelial cells, and thereby induces endocytosis [37,39]. It is notable that *ssa1Δ/Δ* and *als3Δ/Δ* mutants both have defects in their capacity to invade and damage these cells *in vitro*. Furthermore, the host cell invasion defect of the *ssa1Δ/Δ als3Δ/Δ* mutant was not more severe than either parent mutant (data not shown).




Figure 8. Interaction between Ssa1 and Als3. (A) The *ssa1Δ/Δ* mutant has normal amounts of Als3 on its surface. Hyphae of the indicated *C. albicans* strains were stained with a polyclonal anti-Als3 antibody, after which amount of surface exposed Als3 was determined by flow cytometric analysis of 10,000 cells per strain. (B) Normal surface distribution of Als3 on *ssa1Δ/Δ* hyphae. The distribution of Als3 on the surface of hyphae of the indicated *C. albicans* strains was determined by staining the organisms with a polyclonal anti-Als3 antibody (green fluorescence) and counterstaining with an anti-*C. albicans* polyclonal antibody (red fluorescence). (C) Effects of deletion of *ALS3* in the *ssa1Δ/Δ* mutant background. Endothelial cells and FaDu oral epithelial cells were incubated with the indicated strains for 150 min, after which the number of endocytosed and cell-associated organisms were determined. Results are the mean \pm SD of 3 experiments, each performed in triplicate. * $P < 0.01$ compared to the wild-type strain, $^{\dagger}p < 0.005$ compared to the *ssa1Δ/Δ* single mutant. doi:10.1371/journal.ppat.1001181.g008

Δ double mutant was similar that of the *als3Δ/Δ* single mutant. The most likely explanation for this result is that Ssa1 and Als3 bind to same endothelial and epithelial cell surface proteins, either separately or perhaps as part of multiprotein complex. An alternative explanation for these results is that Ssa1 functions as a molecular chaperone that is required for appropriate folding or trafficking of Als3 [47,48,51,52]. Although we cannot completely exclude the possibility that the folding or post-translational modification of Als3 was aberrant in the *ssa1Δ/Δ* mutant, we determined that deletion of *SSA1* had no effect on the amount and surface distribution of Als3 on *C. albicans* hyphae. These results suggest that proper Als3 trafficking can occur in the absence of Ssa1. Furthermore, the localization of Ssa1 on the fungal cell surface and the finding that beads coated with rSsa1 were avidly endocytosed suggests the more likely scenario that Ssa1 and Als3 function cooperatively as invasins.

After *C. albicans* invades host cells, it damages them. Host cell damage requires at least some extent of fungal invasion because blocking invasion by treatment with cytochalasin D or infection with non-invasive mutants of *C. albicans* reduces the extent of host cell damage [27,31,32,50]. Thus, the defective host cell invasion of the *ssa1Δ/Δ* mutant likely contributed to its reduced capacity to damage these cells. Although the exact mechanism by which *C. albicans* causes host cell damage is incompletely understood, it is probable that phospholipases and aspartyl proteases secreted by the organism participate in this process [35,40–46]. Some heat shock proteins can assist with protein secretion [47,48]. However, we found no evidence of impaired phospholipase or protease secretion in the *ssa1Δ/Δ* mutant. While these results indicate that bulk protein secretion is intact in the *ssa1Δ/Δ* mutant, we cannot completely rule out the possibility that there was reduced secretion or activity of a single phospholipase or protease isozyme in this strain.




Figure 9. Relative transcript levels of *SSA1* and *SSA2*. FaDu epithelial cells were infected with the indicated strains of *C. albicans* for 90 min, after which the *C. albicans* RNA was extracted. The relative transcript levels of *SSA1* and *SSA2* were measured by real-time PCR. Results are the mean \pm SD of three biological replicates, each tested in duplicate. * $P < 0.01$ compared to *SSA2* transcript levels. doi:10.1371/journal.ppat.1001181.g009

During both hematogenously disseminated and oropharyngeal infection, *C. albicans* is exposed to significant stress induced by the host. These stresses include exposure to reactive oxygen intermediates and antimicrobial peptides. Furthermore, in the oral cavity, *C. albicans* is likely exposed to toxic secondary metabolites produced by the resident oral flora. In some microbial pathogens, such as *Leishmania spp.* and *Francisella tularensis*, members of the Hsp70 and Hsp100 family of heat shock proteins are required for them to tolerate host-induced stress and cause persistent infection in mice [10–12]. Here, we found that Ssa1 was dispensable for the resistance of *C. albicans* to oxidant, cell wall, and cell membrane stress. We also determined that *ssa1Δ/Δ* mutant cells did not have increased susceptibility to killing by neutrophil-like HL-60 cells. Thus, the reduced virulence of the *ssa1Δ/Δ* mutant was unlikely due to impaired ability to withstand host-induced stress.

Previously we have found that Ssa1 acts as receptor for some antimicrobial proteins such as histatin 5, and that the *ssa1Δ/Δ* mutant is more resistant to some defensins [17]. Thus, one might have predicted that the *ssa1Δ/Δ* mutant would be resistant to neutrophil killing and therefore hypervirulent. However, the




Figure 10. *C. albicans* extracellular phospholipase and protease activities are independent of Ssa1 and Ssa2. The total extracellular phospholipase activity of the indicated strains was determined by growing them on egg yolk agar at room temperature for 5 days and then measuring the width of the zone of precipitation around each colony (top panel). The total extracellular protease activity of the indicated strains was assessed by growing them on BSA agar at 37°C for 5 days and then staining the plates with 0.5% amido black. The extracellular protease activity was proportional to the width of the zone of clearance around the colonies (lower panel). doi:10.1371/journal.ppat.1001181.g010




Figure 11. Ssa1 and Ssa2 are not required for resistance to stress. (A) Susceptibility to oxidant and cell wall stress. Serial 10-fold dilutions of the indicated strains were used to inoculate YPD plates containing menadione, H_2O_2 , Calcofluor white, NaCl, or SDS. Images were obtained after growth at 30°C for 24 h. (B) Susceptibility to leukocyte induced damage. Yeast-phase cells of the indicated strains were incubated for 3 h with HL-60 cells that had been differentiated into neutrophil-like cells, after which the amount of damage to the organisms was determined by an XTT assay. Each strain was tested at an effector to target ratio (E:T) ranging from 5:1 to 1:2. Results are the mean \pm SD of three independent experiments, each performed in triplicate. doi:10.1371/journal.ppat.1001181.g011

ssa1Δ/Δ mutant is not resistant to some toxic products of neutrophils, such as human neutrophil defensin 1 [17], reactive oxygen intermediates, and nitric oxide. Therefore, it is probable that the wild-type susceptibility of the *ssa1Δ/Δ* mutant to these stressors was the reason why it was neither resistant to killing by the neutrophil-like HL-60 cells in vitro nor hypervirulent in mice.

Although induction of host cell endocytosis is an important virulence function of Ssa1, it is possible that this protein may also contribute to *C. albicans* pathogenicity in other ways. For example, in *S. cerevisiae* Ssa1 regulates the transcription of the multidrug resistant gene, *PDR3* [53]. Furthermore, in *Cryptococcus neoformans*, Ssa1 functions as a transcriptional co-activator of laccase, and is required for normal melanin synthesis and virulence [54]. Thus, Ssa1 may either directly or indirectly govern the transcription of genes that also contribute to *C. albicans* pathogenicity. This possibility is currently being investigated.

Materials and Methods

Strains and growth conditions

The *C. albicans* strains used in these studies and their relevant genotypes are listed in Table 1. All strains were maintained on yeast extract/peptone/dextrose (YPD; Qbiogene) agar plates and re-cultured at least monthly from -80°C stock. For use in the experiments, yeast-phase cells of the various strains were grown YPD broth overnight in a rotary shaker at 30°C.

The initial *ssa1Δ/Δ*, *ssa2Δ/Δ*, *ssa1Δ/Δ::SSA1*, and *ssa2Δ/Δ::SSA2* strains were constructed previously [16]. To verify that

Table 1. Strains of *C. albicans* used in this study.

Strain	Genotype				Reference	
SC5314	Wild-type				[55]	
<i>ssa1Δ/Δ</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::FRT</i></u> <i>ssa1::FRT</i>	<u><i>ssa2::FRT::URA3::SSA2</i></u> <i>SSA2</i>		This paper	
<i>ssa1Δ/Δ-URA3</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::FRT</i></u> <i>ssa1::FRT</i>	<u><i>ssa2::FRT</i></u> <i>SSA2</i>	<u><i>rps10::URA3</i></u> <i>RPS10</i>	This paper	
<i>ssa1Δ/SSA1</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::FRT</i></u> <i>ssa1::FRT</i>	<u><i>ssa2::FRT</i></u> <i>SSA2</i>	<u><i>rps10::URA3::SSA1</i></u> <i>RPS10</i>	[16]	
<i>ssa2Δ/Δ</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::FRT::URA3::SSA1</i></u> <i>SSA1</i>	<u><i>ssa2::FRT</i></u> <i>ssa1::FRT</i>		This paper	
<i>Ssa2Δ/SSA2</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa2::FRT</i></u> <i>SSA1</i>	<u><i>ssa2::FRT</i></u> <i>ssa2::FRT</i>	<u><i>rps10::URA3::SSA2</i></u> <i>RPS10</i>	[16]	
CAYF178U	<u><i>ura3Δ::λimm434::URA3-IRO1</i></u> <i>ura3Δ::λimm434</i>	<u><i>als3::ARG4</i></u> <i>als3::HIS1</i>	<u><i>arg4::hisG</i></u> <i>arg4::hisG</i>	<u><i>his1::hisG</i></u> <i>his1::hisG</i>	[68]	
CAQTP178U	<u><i>ura3Δ::λimm434::URA3-IRO1</i></u> <i>ura3Δ::λimm434</i>	<u><i>als3::ARG4::ALS3</i></u> <i>als3::HIS1</i>	<u><i>arg4::hisG</i></u> <i>arg4::hisG</i>	<u><i>his1::hisG</i></u> <i>his1::hisG</i>	[68]	
<i>ssa1Δ/Δ-als3Δ/Δ</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::FRT</i></u> <i>ssa1::FRT</i>	<u><i>ssa2::FRT</i></u> <i>SSA2</i>	<u><i>als3</i></u> <i>als3::NAT1</i>	<u><i>rps10::URA3</i></u> <i>RPS10</i>	This paper
<i>SSA1-GFP</i>	<u><i>ura3Δ::λimm434</i></u> <i>ura3Δ::λimm434</i>	<u><i>ssa1::SSA1-GFP::URA3</i></u> <i>SSA1</i>	<u><i>SSA2</i></u> <i>SSA2</i>		This paper	

doi:10.1371/journal.ppat.1001181.t001

the phenotype of the *ssa1Δ/Δ* mutant was not the result of *URA3* being integrated at the *SSA2* locus, a Ura⁻ strain was selected by growth on 5-fluoroorotic acid [55]. This strain was transformed by the lithium acetate procedure with the CIp10 vector that had been linearized with NcoI [56]. The resulting strain, *ssa1Δ/Δ-URA3* contained *URA3* at the neutral *RPS10* locus [21].

To delete the entire protein coding region of *ALS3* in the *ssa1Δ/Δ* mutant, the PCR-product-directed gene deletion approach was used [57]. Briefly, deletion cassettes containing *ALS3* flanking regions and the *URA3* or *NAT1* selection markers were amplified by PCR with primers GATATTTTGAATATGGAA-ATAAATCGTGCATAAGAAAGTTTGTCTATGCACGTTCA-TACTTCCAAAAATTGTAATACGACTCACTATAGGGC and AAACATATAGAAACAACTAATCAAATTAACAACAC-ACCAAATTGGAGGTAATTAATCATACCGAAAAATAGCT-ATGACCATGATTACGCCA, using pGEM-URA3 [57] and pJK795 [58] as templates, respectively. These PCR products were then used to successively transform the Ura⁻ *ssa1Δ/Δ* strain [57]. The resulting *ssa1Δ/Δ als3Δ/Δ* mutant was plated on 5-fluoroorotic acid to select for a Ura⁻ strain. This strain was transformed with NcoI-linearized CIp10 [56] to re-integrated *URA3* at *RPS10* locus.

Strains containing C-terminal GFP fusions were generated in strain CAF4-2 (wt) by substitution of one *SSA1* allele with an *SSA1-GFP* tagged allele as described previously [16]. Plasmid pGFP-URA3 (PMG1602) was kindly provided by Dr. C. A. Gale, and was used as a template to PCR amplify the transformation cassettes using primers containing sequences flanking the *SSA1* stop codon (67 bp) and homologous sequences from the vector (forward primer: 9 bp glycine linker and 21bp vector sequence; reverse primer: 23 bp vector sequence). PCR products were used to transform parental strains using frozen EZ yeast transformation II kit (Zymo Research) following the manufacturer's protocol and selected on URA3⁻ YNB agar plates. To identify transformants

carrying cassettes correctly integrated into the target gene sequence, genomic DNA was prepared and used as the template for PCR reactions using one primer for annealing within the transformation module and a second primer annealing with the target gene locus outside the altered region. Cells were examined using confocal microscopy and Western blotting with anti-GFP to confirm expression of Ssa1-GFP fusion protein.

Mouse models of disseminated and oropharyngeal candidiasis

The virulence of the various strains was tested in the mouse model of hematogenously disseminated candidiasis as described previously [26]. Briefly, 10 male BALB/c mice (20 g body weight; National Cancer Institute, Bethesda, MD) were infected via the lateral tail vein with 5×10^5 yeast-phase cells of each strain of *C. albicans* in 500 μl of PBS. All inocula were confirmed by quantitative culture. The mice were monitored at least three times daily, and moribund mice were euthanized. To determine the organ fungal burden, 5 to 7 mice were inoculated with each strain as in the survival experiments. After 4 h, 8 h, 1 day, 2 days, 4 days, 7 days, and 14 days of infection, the brain, liver, and one kidney from each mouse were harvested, weighed, homogenized, and quantitatively cultured. The remaining kidneys from the 1 day time point were fixed in zinc-buffered formalin followed by 70% ethanol and then embedded in paraffin. Thin sections were cut and stained with periodic acid-Schiff.

The different strains of *C. albicans* were also tested for virulence in our previously described mouse model of oropharyngeal candidiasis [25,27,59]. Briefly, male BALB/c mice were immunosuppressed with cortisone acetate (225 mg/kg; Sigma-Aldrich) administered subcutaneously on days -1, 1, and 3 relative to infection. For oral inoculation, each mouse was anaesthetized by intraperitoneal injection with ketamine and xylazine (both from Phoenix Pharmaceuticals), after which a calcium alginate swab

(Type 4 Calgiswab; Puritan Medical Products Company LLC) saturated with 10^6 yeast/ml was placed sublingually for 75 min. The mice were subsequently provided food and water ad libitum and then sacrificed after 1, 2 and 5 days of infection. The tongue and adjacent hypoglossal tissue were excised and cut in half. One half was weighed and homogenized for quantitative culture and the other half was processed for histopathological analysis as described above.

Endothelial cells and oral epithelial cells

Endothelial cells were isolated from human umbilical cord veins by the method of Jaff et al. [60] and cultured as described [32] in M-199 medium supplemented with 10% fetal bovine serum and 10% defined bovine calf serum (Gemini Bio-Products), and containing 2 mM L-glutamine with penicillin and streptomycin (Irvine Scientific). Endothelial cells were used at the third passage. The FaDu oral epithelial cell line (American Type Culture Collection) was cultured in Eagle's minimum essential medium with Earle's balanced salt solution (Irvine Scientific) supplemented with 10% fetal bovine serum, 1 mM pyruvic acid, 2 mM L-glutamine, and 0.1 mM nonessential amino acids, with penicillin and streptomycin [25,27]. Both cell types were maintained in a humidified incubator in 5% CO₂ at 37°C.

Host cell damage assay

The extent of endothelial and epithelial cell damage caused by the different strains of *C. albicans* was measured using our previously described ⁵¹Cr release assay [25–27,31,32]. Briefly, endothelial cells or FaDu oral epithelial cells were grown to 95% confluency in 96 well tissue culture plates with detachable wells (Corning) and loaded with 5 µCi/ml Na₂⁵¹CrO₄ (MP Biomedicals) overnight. After removing the unincorporated ⁵¹Cr by rinsing, the cells were then infected with yeast of the various *C. albicans* strains suspended in RPMI 1640 medium (Irvine Scientific). The inoculum was 4×10^4 organisms per well of endothelial cells and 1×10^5 organisms per well of oral epithelial cells. The infected host cells were incubated for 3 h, after which the amount of ⁵¹Cr released into the medium and retained by the cells was determined by γ-counting. Wells containing host cells, but no organism, were processed in parallel to determine the spontaneous release of ⁵¹Cr. After correcting for well-to-well differences in the incorporation of ⁵¹Cr, the percent specific release of ⁵¹Cr was calculated using the following formula: (experimental release - spontaneous release)/(total incorporation - spontaneous release). Experimental release was the amount of ⁵¹Cr released into the medium by cells infected with *C. albicans*. Spontaneous release was the amount of ⁵¹Cr released into the medium by uninfected host cells. Total incorporation was the sum of the amount of ⁵¹Cr released into the medium and remaining in the host cells. Each assay was performed in triplicate on three separate occasions.

Three-dimensional model of the human oral mucosa

The capacity of the various *C. albicans* strains to invade a three dimensional model of the oral mucosa in vitro was determined by our previously described method [34,61]. OKF6/TERT-2 oral epithelial cells were grown on top of a feeder layer of collagen embedded NIH 3T3 cells in 30 mm diameter cell culture inserts (Millipore). They were infected via their apical surface by adding 10^5 *C. albicans* cells in 100 µl of airlift medium (DMEM with 4.5 g/l glucose [Fisher Scientific] and Ham's F-12 medium [Invitrogen] mixed 3:1, and supplemented with 5 µg/ml insulin, 0.4 µg/ml hydrocortisone, 2×10^{-11} M 3,3',5-triiodo-L-thyronine, 1.8×10^{-4} M adenine, 5 µg/ml transferrin, 10^{-10} M cholera toxin,

2 mM L-glutamine; 5% FBS, and penicillin–streptomycin). For histopathology, the cultures were fixed after 2 days with 10% formaldehyde in PBS and embedded in paraffin. Thin sections were stained with periodic acid-Schiff and then evaluated by light microscopy. The extent of epithelial cell damage was quantified after 2 days of infection by measuring the accumulation of LDH in the medium using the CytoTox-96 assay (Promega).

Adherence and endocytosis assay

The capacity of the various *C. albicans* strains to adhere to and be endocytosed by endothelial and FaDu epithelial cells was quantified using our previously described differential fluorescence assay [25,27,32,36]. The host cells were grown to 95% confluency on fibronectin-coated glass coverslips in a 24-well tissue culture plate. Each coverslip was infected with 10^5 organisms in RPMI 1640 medium. After a 90 or 150 min incubation, the medium was aspirated, non-adherent organisms were removed by rinsing the coverslips with Hank's balanced salt solution (HBSS; Irvine Scientific), and cells were fixed with 3% paraformaldehyde. The adherent and non-internalized portions of the organisms were stained with rabbit anti-*C. albicans* antiserum (Biodesign International) conjugated with Alexa 594 (Molecular Probes). Next, the host cells were permeabilized with 1% Triton X-100 for 15 min, and then all of the *C. albicans* cells were stained with rabbit anti-*C. albicans* antiserum conjugated with Alexa 488 (Molecular Probes). The coverslips were mounted inverted on a microscope slide and organisms were viewed under epifluorescence. The number of endocytosed organisms was determined by subtracting the number of non-endocytosed organisms (which fluoresced red) from the number of cell-associated organisms (endocytosed plus non-endocytosed organisms, which fluoresced green). Organisms that were partially internalized were considered to be endocytosed. At least 100 organisms were counted per coverslip, and each experiment was repeated in triplicate, at least three times. Results were expressed as the number of endocytosed and cell-associated organisms per high powered field.

C. albicans binding to endothelial cell N-cadherin and epithelial cell E-cadherin

The capacity of each strain to bind to N-cadherin and E-cadherin was determined using our affinity purification procedure as outlined previously [36,37]. Germ tubes were prepared by incubating yeast-phase cells for 90 min at 37°C in 150 mm diameter Petri dishes containing RPMI 1640 medium buffered to pH 7.5 with 150 mM HEPES. The resulting germ tubes were removed by scraping and rinsed once with PBS containing calcium and magnesium. The germ tubes (2×10^8 cells) were incubated for 1 h on ice with 250 µg of cell membrane proteins of endothelial cells or FaDu epithelial cells in PBS with calcium and magnesium containing 1.5% octyl-glucopyranoside and protease inhibitors. The unbound proteins were removed by rinsing in the same buffer, after which the proteins that remained bound to the germ tube were eluted with 6 M urea. The eluted proteins were separated by SDS-PAGE. N-cadherin and E-cadherin were detected by immunoblotting with anti-N-cadherin murine monoclonal antibody (clone 32, Transduction Laboratories), and an anti-E-cadherin murine monoclonal antibody (clone HECD 1), respectively.

Host cell interactions of latex beads coated with rSsa1, rSsa2, or BSA

Recombinant Ssa1 (rSsa1) and rSsa2 were produced in *S. cerevisiae* as outlined previously [16]. Latex beads were coated with these proteins or biotinylated BSA as described [37,62]. Briefly,

5×10^7 fluorescent, yellow-green, amine-modified, 2.0 μm diameter polystyrene latex beads (Sigma-Aldrich) were washed with PBS followed by coupling buffer (0.2 M Na_2HCO_3 [pH 8.5], 0.5 M NaCl). The beads were then incubated with rSsa1, rSsa2 (0.5 mg/ml) or coupling buffer at 37°C for 30 min. The beads that had been coated with Ssa proteins were incubated with 1% rabbit serum, while the control beads were incubated with biotinylated BSA (1 mg/ml). All beads were sonicated briefly, blocked with unlabeled BSA and rinsed PBS-BSA. They were suspended in PBS containing 2 mg BSA per ml. Binding of rSsa1 or rSsa2 to the beads was verified by indirect immunofluorescence with an anti-Xpress monoclonal antibody (Invitrogen) directed against the Xpress leader sequence of these recombinant proteins. Binding of biotinylated BSA to the beads was confirmed using Alexa 568 conjugated streptavidin (Invitrogen). The interactions of these beads with endothelial and FaDu epithelial cells were determined by the differential fluorescent assay described above.

Real-time PCR detection of *SSA1* and *SSA2* transcript levels

The relative transcript levels of *SSA1* and *SSA2* in hyphae of the various strains of *C. albicans* were determined by our previously described method [63]. Yeast cells were suspended in RPMI 1640 medium and added to FaDu epithelial cells at a final concentration of 5×10^5 cells/cm². After a 90 min incubation, the non-adherent organisms were removed by rinsing with ice-cold distilled water. Next, ice-cold DEPC-treated water was added and the *C. albicans* and epithelial cells were removed with a cell scraper. The mixture was vortexed for 30 sec to lyse the epithelial cells and then the organisms were collected by a brief centrifugation at 4°C. These organisms were suspended in TES buffer (10 mM Tris, 10 mM EDTA, 0.5% SDS), and then snap frozen in liquid nitrogen. The total time from rinsing the organisms to freezing them in liquid nitrogen was less than 5 min. At a later time, the cells were thawed on ice and the fungal RNA was extracted by the hot phenol method [64]. For real-time PCR, the *C. albicans* RNA was treated with DNase I (Ambion), after which cDNA was synthesized using MMLV reverse transcriptase (Ambion). Quantitative real-time PCR was carried out using the SYBR green PCR kit (Applied Biosystems) and an ABI 7000 Real-Time PCR System (Applied Biosystems) following the manufacturer's protocol. The results were analyzed by the $2^{-\Delta\Delta\text{CT}}$ method [65] using *ACT1* as the endogenous control. The *SSA1* and *SSA2* transcript levels were determined in three biological replicates, each tested in duplicate.

Assessment of surface expression of Als3 and Ssa1

Flow cytometry was used to quantify the amount of Als3 on the surface of the various strains using a slight modification of our previously described method [37]. Briefly, 3×10^{10} yeast-phase cells in 15 ml RPMI 1640 medium buffered to pH 7.5 with 10 mM HEPES were added to 100 mm diameter Petri dishes and incubated in 5% CO_2 at 37°C. After 90 min, the resulting germ tubes were scraped from the dishes with a cell scraper, fixed with 3% paraformaldehyde, blocked with 1% goat serum, and then stained with a rabbit polyclonal anti-Als3 antiserum [37] followed by an Alexa 488 conjugated goat anti-rabbit antibody (Molecular Probes). An *als3* Δ/Δ mutant was included in these experiments as a negative control. The amount of Als3 surface staining on 10^4 germ tubes per strain was analyzed by flow cytometry.

To analyze the surface distribution of Als3 on the various strains, 10^5 yeast-phase cells in 1 ml RPMI 1640 medium buffered to pH 7.5 with 10 mM HEPES were added to 12 mm diameter glass cover slips in a 24-well tissue culture plate. After a 90 min incubation in 5% CO_2 at 37°C, the resulting germ tubes were

fixed with 3% paraformaldehyde, and blocked with 1% goat serum, and then stained with the rabbit polyclonal anti-Als3 antiserum followed by an Alexa 488 conjugated goat anti-rabbit antibody. Next, the germ tubes were counter stained with the Alexa 594 conjugated anti-*Candida* antibody to label the cell surface. The coverslips were mounted inverted on microscope slides and imaged by confocal microscopy. Stacked images were acquired along the z-axis and then combined for the final images.

The surface expression of Ssa1 on *C. albicans* expressing Ssa1-GFP was determined by a similar process, except that FaDu oral epithelial cells were grown on the coverslips prior to adding the organisms and the anti-Als3 antiserum was omitted.

Screening for extracellular phospholipase and protease activities

The *C. albicans* strains were screened for phospholipase activity by the method of Samaranayake et al. [49]. Briefly, egg yolk agar consisting of Sabouraud's dextrose agar, NaCl, CaCl_2 and egg yolk emulsion was prepared. A 10 μl of suspension of 10^7 yeast-phase cells per ml in 10 mM sodium phosphate buffer was plated on the agar and incubated at room temperature for 5 days. The phospholipase activity of each strain was determined by measuring the width of zone of precipitation around the colony.

The total extracellular protease activity of the various strains was assessed using BSA agar by the method of Ruchel et al. [66]. BSA agar (0.2% BSA, 1.17% yeast carbon base, and 0.01% yeast extract, pH 5) was spot inoculated with the various *C. albicans* strains and then incubated at 37°C. After 5 days the agar was stained with 0.5% amido black and the width of the zone of clearance around each colony was measured.

Susceptibility to stressors

The susceptibility of the various *C. albicans* strains to stressors were tested by spotting dilutions of 10^4 to 10^1 yeast-phase cells in a total volume of 7 μl onto YPD agar plates containing menadione (12.5 and 25 μM); H_2O_2 (1, 2 and 4 mM), Calcafluor white (25 and 50 μM), NaCl (1 and 2 M) and SDS (0.002%). The plates were incubated at 30°C for 24 hours and then imaged.

The susceptibility of the different strains to damage by a neutrophil-like cell line was determined by the 2,3-bis (2-methoxy-4-nitro-5-sulphophenyl)-5-[(phenylamino) carbonyl]-2H-tetrazolium hydroxide (XTT) assay, as previously described [67]. Briefly, HL-60 cells were cultured in RPMI 1640 medium containing 10% fetal bovine serum and 25 mM HEPES. They were induced to differentiate into neutrophil-like cells by exposure to 1.25% dimethyl sulfoxide for 7 days. For the damage assay, *C. albicans* yeast suspended in Dulbecco's Modified Eagle Medium containing 10% fetal bovine serum were added to 96-well plates at concentrations ranging from 10^5 to 2×10^4 cells/well. There was a linear relationship between viable cell number and colorimetric signal (XTT activity) in this concentration range with all three strains (not shown). HL-60 cells were added to the *C. albicans* cells at effector to target cell ratios (E:T) ranging from 5:1 to 1:2. After incubation at 37°C in 5% CO_2 for 3 hours, the medium was aspirated and the HL-60 cells were lysed with sterile H_2O . To each well was added 100 μl of a mixture of XTT and coenzyme Q0 (0.25 mg/ml XTT and 40 μg /ml coenzyme Q0), after which the plate was incubated at 37°C in 5% CO_2 for 2 h. Supernatants were transferred into new plates, and optical densities (OD) were measured by an Opsys Microplate Reader (Thermo Labsystems) at 450–490 nm, with a 630 nm reference filter. Antifungal activity was calculated according to the following formula: %fungal damage = $(1 - x/n) \times 100$, where x is the OD450 of experimental wells (*C. albicans* with effectors) and n is the OD450 of control wells

(*C. albicans* only). Each experiment was performed in triplicate and repeated 3 times.

Statistics

The results of the survival experiments were analyzed with the Log-Rank Test, and the organ fungal burden data were analyzed using the Log Rank Test. The results of the in vitro experiments were analyzed using the Student's T test. *P* values ≤ 0.05 were considered to be significant.

Ethics statement

All experiments were approved by the Los Angeles Biomedical Research Institute Animal Care and Use Committee and as outlined in the Guide for the Care and Use of Laboratory Animals of National Institutes of Health. The collection of umbilical cords for the harvesting of endothelial cells used in these studies was approved by the Institutional Review Board of the Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center.

Supporting Information

Figure S1 The chromosomal location of *URA3* does not influence the virulence or host cell interactions of the *ssa1Δ/Δ* mutant. (A). Survival of mice infected with 5×10^5 yeast cells of the indicated strains of *C. albicans*. Each strain was used to inoculate 10 mice. (B) Endothelial cells and FaDu oral epithelial cells were incubated with the indicated strains for 90 min, after which the

number of endocytosed and cell-associated organisms were determined. Results are the mean \pm SD of 3 experiments, each performed in triplicate.

Found at: doi:10.1371/journal.ppat.1001181.s001 (0.45 MB TIF)

Figure S2 Localization of Ssa1 on *C. albicans* yeast. *C. albicans* yeast expressing an Ssa1-GFP fusion protein were stained with an Alexa 594-conjugated anti-*C. albicans* polyclonal antibody to label the cell surface and then imaged by confocal microscopy. (A–C) Images of Ssa1-GFP (A) and the fluorescent-labeled anti-*C. albicans* antibody (B). The merged image is shown in (C). (D) Graphs of fluorescent intensity at different cross sections of the yeast in panel (C). The green lines indicate the fluorescent intensity of the Ssa1-GFP and the red lines indicate the fluorescent intensity of the fluorescent-labeled anti-*C. albicans* antibody.

Found at: doi:10.1371/journal.ppat.1001181.s002 (1.38 MB TIF)

Acknowledgments

We thank Damien Tahbaz for help with tissue culture and perinatal nurses at the Harbor-UCLA Medical Center Pediatric Clinical Research Center for collection of umbilical cords.

Author Contributions

Conceived and designed the experiments: JNS ADB ME SGF. Performed the experiments: JNS NVS QTP JSB HK AT YL. Analyzed the data: JNS ADB ME SGF. Wrote the paper: JNS ADB ME SGF.

References

- Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, et al. (2004) Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. *Clin Infect Dis* 39: 309–317.
- Filler SG (2006) *Candida*-host cell receptor-ligand interactions. *Curr Opin Microbiol* 9: 333–339.
- Grubb SE, Murdoch C, Sudbery PE, Saville SP, Lopez-Ribot JL, et al. (2008) *Candida albicans*-endothelial cell interactions: a key step in the pathogenesis of systemic candidiasis. *Infect Immun* 76: 4370–4377.
- Walker LA, Maccallum DM, Bertram G, Gow NA, Odds FC, et al. (2009) Genome-wide analysis of *Candida albicans* gene expression patterns during infection of the mammalian kidney. *Fungal Genet Biol* 46: 210–219.
- Rodaki A, Bohovych IM, Enjalbert B, Young T, Odds FC, et al. (2009) Glucose promotes stress resistance in the fungal pathogen *Candida albicans*. *Mol Biol Cell* 20: 4845–4855.
- Yamaguchi H, Osaki T, Taguchi H, Hanawa T, Yamamoto T, et al. (1996) Flow cytometric analysis of the heat shock protein 60 expressed on the cell surface of *Helicobacter pylori*. *J Med Microbiol* 45: 270–277.
- Hartmann E, Lingwood C (1997) Brief heat shock treatment induces a long-lasting alteration in the glycolipid receptor binding specificity and growth rate of *Haemophilus influenzae*. *Infect Immun* 65: 1729–1733.
- Ratnakar P, Rao SP, Catanzaro A (1996) Isolation and characterization of a 70 kDa protein from *Mycobacterium avium*. *Microb Pathog* 21: 471–486.
- Long KH, Gomez FJ, Morris RE, Newman SL (2003) Identification of heat shock protein 60 as the ligand on *Histoplasma capsulatum* that mediates binding to CD18 receptors on human macrophages. *J Immunol* 170: 487–494.
- Hubel A, Krobitsch S, Harauf A, Clos J (1997) *Leishmania major* Hsp100 is required chiefly in the mammalian stage of the parasite. *Mol Cell Biol* 17: 5987–5995.
- Folgueira C, Carrion J, Moreno J, Saugar JM, Canavate C, et al. (2008) Effects of the disruption of the HSP70-II gene on the growth, morphology, and virulence of *Leishmania infantum* promastigotes. *Int Microbiol* 11: 81–89.
- Meibom KL, Dubail I, Dupuis M, Barel M, Lencio J, et al. (2008) The heat-shock protein ClpB of *Francisella tularensis* is involved in stress tolerance and is required for multiplication in target organs of infected mice. *Mol Microbiol* 67: 1384–1401.
- Urban C, Sohn K, Lottspeich F, Brunner H, Rupp S (2003) Identification of cell surface determinants in *Candida albicans* reveals Tsalp, a protein differentially localized in the cell. *FEBS Lett* 544: 228–235.
- Lopez-Ribot JL, Alloush HM, Masten BJ, Chaffin WL (1996) Evidence for presence in the cell wall of *Candida albicans* of a protein related to the hsp70 family. *Infect Immun* 64: 3333–3340.
- Li XS, Reddy MS, Baev D, Edgerton M (2003) *Candida albicans* Ssa1/2p is the cell envelope binding protein for human salivary histatin 5. *J Biol Chem* 278: 28553–28561.
- Li XS, Sun JN, Okamoto-Shibayama K, Edgerton M (2006) *Candida albicans* cell wall Ssa proteins bind and facilitate import of salivary histatin 5 required for toxicity. *J Biol Chem* 281: 22453–22463.
- Vylkova S, Li XS, Berner JC, Edgerton M (2006) Distinct antifungal mechanisms: beta-defensins require *Candida albicans* Ssa1 protein, while Trk1p mediates activity of cysteine-free cationic peptides. *Antimicrob Agents Chemother* 50: 324–331.
- Sundstrom P, Cutler JE, Staab JF (2002) Reevaluation of the role of *HWPI* in systemic candidiasis by use of *Candida albicans* strains with selectable marker *URA3* targeted to the *ENO1* locus. *Infect Immun* 70: 3281–3283.
- Cheng S, Nguyen MH, Zhang Z, Jia H, Handfield M, et al. (2003) Evaluation of the roles of four *Candida albicans* genes in virulence by using gene disruption strains that express *URA3* from the native locus. *Infect Immun* 71: 6101–6103.
- Staab JF, Sundstrom P (2003) *URA3* as a selectable marker for disruption and virulence assessment of *Candida albicans* genes. *Trends Microbiol* 11: 69–73.
- Brand A, MacCallum DM, Brown AJ, Gow NA, Odds FC (2004) Ectopic Expression of *URA3* can influence the virulence phenotypes and proteome of *Candida albicans* but can be overcome by targeted reintegration of *URA3* at the *RPS10* locus. *Eukaryot Cell* 3: 900–909.
- Eversole LR, Reichart PA, Ficarra G, Schmidt-Westhausen A, Romagnoli P, et al. (1997) Oral keratinocyte immune responses in HIV-associated candidiasis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 84: 372–380.
- Montes LF, Wilborn WH (1968) Ultrastructural features of host-parasite relationship in oral candidiasis. *J Bacteriol* 96: 1349–1356.
- Cawson RA, Rajasingham KC (1972) Ultrastructural features of the invasive phase of *Candida albicans*. *Br J Dermatol* 87: 435–443.
- Chiang LY, Sheppard DC, Bruno VM, Mitchell AP, Edwards JE, Jr., et al. (2007) *Candida albicans* protein kinase CK2 governs virulence during oropharyngeal candidiasis. *Cell Microbiol* 9: 233–245.
- Sanchez AA, Johnston DA, Myers C, Edwards JE, Jr., Mitchell AP, et al. (2004) Relationship between *Candida albicans* virulence during experimental hematogenously disseminated infection and endothelial cell damage in vitro. *Infect Immun* 72: 598–601.
- Park H, Myers CL, Sheppard DC, Phan QT, Sanchez AA, et al. (2005) Role of the fungal Ras-protein kinase A pathway in governing epithelial cell interactions during oropharyngeal candidiasis. *Cell Microbiol* 7: 499–510.
- Chopra J, Joist JH, Webster RO (1987) Loss of ^{51}Cr chromium, lactate dehydrogenase, and ^{111}In indium as indicators of endothelial cell injury. *Lab Invest* 57: 578–584.
- Rotrosen D, Edwards JE, Jr., Gibson TR, Moore JC, Cohen AH, et al. (1985) Adherence of *Candida* to cultured vascular endothelial cells: mechanisms of attachment and endothelial cell penetration. *J Infect Dis* 152: 1264–1274.
- Filler SG, Ibe BO, Luckett PM, Raj JU, Edwards JE, Jr. (1991) *Candida albicans* stimulates endothelial cell eicosanoid production. *J Infect Dis* 164: 928–935.

31. Filler SG, Swerdloff JN, Hobbs C, Luckett PM (1995) Penetration and damage of endothelial cells by *Candida albicans*. *Infect Immun* 63: 976–983.
32. Phan QT, Belanger PH, Filler SG (2000) Role of hyphal formation in interactions of *Candida albicans* with endothelial cells. *Infect Immun* 68: 3485–3490.
33. Zakikhany K, Naglik JR, Schmidt-Westhausen A, Holland G, Schaller M, et al. (2007) In vivo transcript profiling of *Candida albicans* identifies a gene essential for interepithelial dissemination. *Cell Microbiol* 9: 2938–2954.
34. Dongari-Bagtzoglou A, Kashleva H (2006) Development of a novel three-dimensional in vitro model of oral *Candida* infection. *Microb Pathog* 40: 271–278.
35. Dalle F, Wachtler B, Coralie L, Holland G, Bannert N, et al. (2010) Cellular interactions of *Candida albicans* with human oral epithelial cells and enterocytes. *Cell Microbiol* 12: 248–271.
36. Phan QT, Fratti RA, Prasadarao NV, Edwards JE, Jr., Filler SG (2005) N-cadherin mediates endocytosis of *Candida albicans* by endothelial cells. *J Biol Chem* 280: 10455–10461.
37. Phan QT, Myers CL, Fu Y, Sheppard DC, Yeaman MR, et al. (2007) Als3 is a *Candida albicans* invasin that binds to cadherins and induces endocytosis by host cells. *PLoS Biol* 5: e64.
38. Zhao X, Oh SH, Cheng G, Green CB, Nuessen JA, et al. (2004) *ALS3* and *ALS8* represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology* 150: 2415–2428.
39. Moreno-Ruiz E, Galan-Diez M, Zhu W, Fernandez-Ruiz E, d'Enfert C, et al. (2009) *Candida albicans* internalization by host cells is mediated by a clathrin-dependent mechanism. *Cell Microbiol* 11: 1179–1189.
40. Leidich SD, Ibrahim AS, Fu Y, Koul A, Jessup C, et al. (1998) Cloning and disruption of *caPLB1*, a phospholipase B gene involved in the pathogenicity of *Candida albicans*. *J Biol Chem* 273: 26078–26086.
41. Ibrahim AS, Mirbod F, Filler SG, Banno Y, Cole GT, et al. (1995) Evidence implicating phospholipase as a virulence factor of *Candida albicans*. *Infect Immun* 63: 1993–1998.
42. Theiss S, Ishdorj G, Brenot A, Kretschmar M, Lan CY, et al. (2006) Inactivation of the phospholipase B gene *PLB5* in wild-type *Candida albicans* reduces cell-associated phospholipase A2 activity and attenuates virulence. *Int J Med Microbiol* 296: 405–420.
43. De Bernardis F, Arancia S, Morelli L, Hube B, Sanglard D, et al. (1999) Evidence that members of the secretory aspartyl proteinase gene family, in particular *SAP2*, are virulence factors for *Candida vaginitis*. *J Infect Dis* 179: 201–208.
44. Hube B, Sanglard D, Odds FC, Hess D, Monod M, et al. (1997) Disruption of each of the secreted aspartyl proteinase genes *SAP1*, *SAP2*, and *SAP3* of *Candida albicans* attenuates virulence. *Infect Immun* 65: 3529–3538.
45. Sanglard D, Hube B, Monod M, Odds FC, Gow NA (1997) A triple deletion of the secreted aspartyl proteinase genes *SAP4*, *SAP5*, and *SAP6* of *Candida albicans* causes attenuated virulence. *Infect Immun* 65: 3539–3546.
46. Schaller M, Korting HC, Schafer W, Bastert J, Chen W, et al. (1999) Secreted aspartic proteinase (Sap) activity contributes to tissue damage in a model of human oral candidosis. *Mol Microbiol* 34: 169–180.
47. Rakestraw A, Wittrup KD (2006) Contrasting secretory processing of simultaneously expressed heterologous proteins in *Saccharomyces cerevisiae*. *Biotechnol Bioeng* 93: 896–905.
48. Holkeri H, Paunola E, Jamsa E, Makarow M (1998) Dissection of the translocation and chaperoning functions of yeast BiP/Kar2p in vivo. *J Cell Sci* 111(Pt 6): 749–757.
49. Samaranyake LP, Raeside JM, MacFarlane TW (1984) Factors affecting the phospholipase activity of *Candida* species in vitro. *Sabouraudia* 22: 201–207.
50. Nobile CJ, Solis N, Myers CL, Fay AJ, Deneault JS, et al. (2008) *Candida albicans* transcription factor Rim101 mediates pathogenic interactions through cell wall functions. *Cell Microbiol* 10: 2180–2196.
51. Yam AY, Albanese V, Lin HT, Frydman J (2005) Hsp110 cooperates with different cytosolic HSP70 systems in a pathway for de novo folding. *J Biol Chem* 280: 41252–41261.
52. Kim S, Schilke B, Craig EA, Horwich AL (1998) Folding in vivo of a newly translated yeast cytosolic enzyme is mediated by the SSA class of cytosolic yeast Hsp70 proteins. *Proc Natl Acad Sci U S A* 95: 12860–12865.
53. Shahi P, Gulshan K, Moye-Rowley WS (2007) Negative transcriptional regulation of multidrug resistance gene expression by an Hsp70 protein. *J Biol Chem* 282: 26822–26831.
54. Zhang S, Hacham M, Panepinto J, Hu G, Shin S, et al. (2006) The Hsp70 member, Ssa1, acts as a DNA-binding transcriptional co-activator of laccase in *Cryptococcus neoformans*. *Mol Microbiol* 62: 1090–1101.
55. Fonzi WA, Irwin MY (1993) Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics* 134: 717–728.
56. Murad AM, Lee PR, Broadbent ID, Barelle CJ, Brown AJ (2000) Clp10, an efficient and convenient integrating vector for *Candida albicans*. *Yeast* 16: 325–327.
57. Wilson RB, Davis D, Mitchell AP (1999) Rapid hypothesis testing with *Candida albicans* through gene disruption with short homology regions. *J Bacteriol* 181: 1868–1874.
58. Shen J, Guo W, Kohler JR (2005) CaNAT1, a heterologous dominant selectable marker for transformation of *Candida albicans* and other pathogenic *Candida* species. *Infect Immun* 73: 1239–1242.
59. Kamai Y, Kubota M, Hosokawa T, Fukuoaka T, Filler SG (2001) New model of oropharyngeal candidiasis in mice. *Antimicrob Agents Chemother* 45: 3195–3197.
60. Jaffe EA, Nachman RL, Becker CG, Minick CR (1973) Culture of human endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. *J Clin Invest* 52: 2745–2756.
61. Villar CC, Kashleva H, Nobile CJ, Mitchell AP, Dongari-Bagtzoglou A (2007) Mucosal tissue invasion by *Candida albicans* is associated with E-cadherin degradation, mediated by transcription factor Rim101p and protease Sap5p. *Infect Immun* 75: 2126–2135.
62. Dersch P, Isberg RR (1999) A region of the *Yersinia pseudotuberculosis* invasin protein enhances integrin-mediated uptake into mammalian cells and promotes self-association. *Embo J* 18: 1199–1213.
63. Park H, Liu Y, Solis N, Spotkov J, Hamaker J, et al. (2009) Transcriptional responses of *Candida albicans* to epithelial and endothelial cells. *Eukaryot Cell* 8: 1498–1510.
64. Kohrer K, Domdey H (1991) Preparation of high molecular weight RNA. *Methods Enzymol* 194: 398–405.
65. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC_T} Method. *Methods* 25: 402–408.
66. Ruchel R, Tegeler R, Trost M (1982) A comparison of secretory proteinases from different strains of *Candida albicans*. *Sabouraudia* 20: 233–244.
67. Dongari-Bagtzoglou A, Villar CC, Kashleva H (2005) *Candida albicans*-infected oral epithelial cells augment the anti-fungal activity of human neutrophils in vitro. *Med Mycol* 43: 545–549.
68. Nobile CJ, Andes DR, Nett JE, Smith EJ, Yue F, et al. (2006) Critical role of Bcr1-dependent adhesins in *C. albicans* biofilm formation in vitro and in vivo. *PLoS Pathog* 2: e63.